

## Combination of the Chimney Technique and an iliac-branched device for the repair of a failed EVAR

Petroula Nana<sup>1</sup>, George Kouvelos<sup>1</sup>, Konstantinos Batzalexis<sup>1</sup>, Konstantinos Spanos<sup>1</sup>, Katerina Koutsothymiou<sup>2</sup>, Eleni Arnaoutoglou<sup>2</sup>, Miltiadis Matsagkas<sup>1</sup>

<sup>1</sup>Department of Vascular Surgery, Faculty of Medicine, School of Health Sciences, University of Thessaly, Larissa, Greece

<sup>2</sup>Department of Anesthesiology, Faculty of Medicine, School of Health Sciences, University of Thessaly, Larissa, Greece

### Abstract:

The era to confront the long-term complications of endovascular aortic aneurysm repair (EVAR) has arisen, while a complex endovascular approach may be usually demanded in these cases. We present an 83-year old male patient, with a failed EVAR, applied 5 years ago. Computed tomography angiography revealed a large increase in aneurysm diameter (93mm) due to endoleak type Ia (ET Ia) and bilateral type Ib, caused by proximal neck dilatation and bilateral common iliac artery enlargement. An aortic graft extension, using a thoracic endograft, and 3 parallel grafts for both renal and superior mesenteric arteries were used to deal with the ET Ia. An iliac branched device for the preservation of the left internal iliac artery as well as coiling of the right one and extension to the external iliac were used to treat ET Ib. The patient was discharged the 6th post-operative day and the post-operative 1st year of follow-up was uneventful.

### INTRODUCTION

The number of cases with failed endovascular aneurysm repair (EVAR) is suspected to increase over the upcoming years.<sup>1</sup> During follow-up, proximal neck dilatation may be present in almost 25% of patients previously treated with EVAR. Neck aneurysmal evolution is related to worse clinical outcomes, due to a higher incidence of type Ia endoleak (ET Ia), migration, and reinterventions needed.<sup>2</sup> In the iliac sealing zone, iliac diameter may exceed graft diameter in mid-term follow-up in up to one third of patients and may be associated with endograft retraction and ET Ib evolution.<sup>3</sup> In such cases, the application of different available treatment modalities may have an important impact in patients' outcomes.<sup>1</sup> Herein, we report a case of an 83-year-old male, with a failed EVAR, treated with a combination of the chimney technique and iliac-branched device. This report has been approved by the Ethics Committee of the Hospital.

### CASE REPORT

An 83-year old male, with a medical history of EVAR, tobacco use, hypertension, dyslipidemia, coronary artery disease with decreased ejection fraction (EF 35%) and chronic obstructive

pulmonary disease presented after 5 years loss of follow-up. He has been treated for an infra-renal abdominal aortic aneurysm of 85mm, associated to iliac aneurysms of 21-23mm, using a bifurcated endoprosthesis with a long main body (Treovance, Bolton, Sunrise, USA) (Figure 1).



Author for correspondence:

**Miltiadis Matsagkas, MD, PhD, FEBVS**



Professor of Vascular Surgery, Department of Vascular Surgery, Medical School, University of Thessaly, Mezourlo, Larissa, Greece

Tel: +30 2413501739

E-mail: mil mats@gmail.com


ISSN 1106-7237/ 2019 Hellenic Society of Vascular and Endovascular Surgery Published by Rotonda Publications  
All rights reserved. <https://www.heljves.com>

A month later, the aneurysm sac had already begun to shrink at 79mm. The patient was in compliant to the surveillance protocol and presented 5 years afterwards due to a diffused abdominal pain. Computed tomography angiography revealed an ET Ia due to neck dilatation without graft migration, sac expansion at 93mm and bilateral ET Ib due to iliac aneurysms. (Figure 2 & 3)




Extreme aortic and iliac tortuosity and lack of time precluded any treatment with fenestrated or branched devices (Figure 4). Considering the complex anatomy, patient's general sta-

tus and the emergent character of the case, a proximal extension using a short thoracic endograft with 3 parallel grafts for visceral vessels (two renals and a superior mesenteric artery (SMA)), was decided to achieve sealing in the upper part. Taking into account the presence of the previous long main body endograft (100mm) and the large supra-renal diameter, the decision to use a thoracic graft of 38mm in diameter seemed rational. A decision to use an iliac branched device for the preservation of the left internal iliac artery (IIA) and a coil embolization for the right IIA in conjunction to an iliac limb extension to the right external iliac artery (EIA) was also made.



Access was achieved using both axillary and femoral arteries. Due to the high tortuosity, via the left common femoral artery, a crossover wire was extracted to the right axillary artery using a snare. The left IIA was catheterized from above and the iliac-branched device was deployed (Iliac Branched Endoprosthesis, Gore, Newark, USA). A self-expanding covered stent (13x50mm, Viabahn, Gore, Newark, USA) was applied as a bridging stent into the left IIA (Figure 5).







Parallel covered stents (Be-Graft, Bentley, Innomed, Germany) were deployed into both renal arteries (RA) and SMA to achieve perfusion of kidneys and bowel. All three splanchnic vessels were catheterized from above. The thoracic endograft was inserted from the right femoral artery and deployed up to the celiac trunk (38x38x100, Relay, Bolton, Sunrise, USA). All target vessel stents were relined using self-expanding stents to preserve patency (E-luminexx, Bard, Covington, USA) (Figure 6). Subsequently, using the left axillary artery, the right IIA was catheterized and embolized using coils (10x20mm, Bolton Medical, Sunrise, USA). Finally, an additional limb extension, down to the EIA, was used to seal the iliac aneurysm (Excluder, 16x14.5x120mm, W. L. Gore & Associates, Newark, USA). The completion angiography showed no endoleak and all stents were patent. The operation duration was 240 min, contrast used was 180ml and radiation exposure (dose area product) was 606mGy (102 min).



Pre-discharge CTA revealed the presence of a small gutter endoleak (Figure 7). The patient was discharged the 6<sup>th</sup> post-op-

erative day without any complication recorded. First-month follow-up with CTA confirmed the resolution of the gutter endoleak and a sac diameter decreased at 86mm. Follow-up at 6 and 12 months revealed no complication, with complete sac exclusion and patent visceral and IIA stents, while there is a moderate shrinkage of the sac to 79mm (Figure 8).



**DISCUSSION**

Proximal neck diameter increases after EVAR, mainly due to the continuous evolution of the aneurysmal disease.<sup>4</sup> In the majority of patients, the aortic neck is suspected to present an increase of  $5.9 \pm 9.3\%$  during the early follow-up.<sup>4</sup> In this case, the patient was treated for an intact aneurysm of

83mm. Sac diameter was begun to decreased during the early post-operative period. However, due to patient's non-compliance with the follow-up protocol, no further information was available for the next 5 years. At presentation, an important neck dilatation affected proximal sealing and contributed to the development of an ET Ia. Nevertheless, aneurysmal evolution may also affect distal sealing and evoke a concomitant common iliac artery expansion.<sup>5</sup> In this patient, iliac dilatation resulted in a bilateral ET Ib which further affected sac remodeling. Iliac dilatation and endograft migration may be detected in patients treated with EVAR, especially when the initial iliac diameter exceeds 24mm.<sup>3</sup>

Current guidelines recommend the use of endovascular means for the management of endoleaks as the first line treatment choice.<sup>6</sup> Type I endoleaks (ET Ia and b) should be treated immediately; using coil or glue embolization, proximal extension with a chimney or conversion to a fenestrated or branched endografts in case of proximal leak or distal extension of the iliac limb with or without IIA preservation; in selected cases down to the EIA.<sup>7</sup> Especially, in this elder patient, an endovascular approach would be the safest solution due to his multiple comorbidities. In high comorbid elder patients, endovascular treatment is associated with acceptable mortality while re-intervention rate is comparable to younger patients.<sup>8</sup>

Neck degeneration and graft migration after EVAR may set the indication for proximal extension using different endovascular approaches; as fenestrated and branched devices, parallel grafts (chimney technique) and endo-staplers.<sup>1,9,10</sup> In any case, the endovascular management of proximal neck dilatation is associated with lower 30-day and aneurysm related mortality.<sup>1</sup> After a careful evaluation of the pre-operative CTA and due to the symptomatic aneurysm expansion, a proximal extension using the parallel grafts technique was used in this patient. Important neck dilatation, proximal aortic diameter and tortuosity precluded any other endovascular technique than chimney EVAR. This off-the-shelf approach permitted the immediate and effective treatment of this patient who presented an important sac increase associated to atypical abdominal symptoms.

Distally, an iliac branched device was used to preserve left IIA flow while contra-laterally, a coil embolization of the IIA associated with an EIA extension were used. No complications were recorded. The adverse anatomy, the longer operational time as well as the higher cost hampered the use of bilateral iliac branched devices while patient's age and comorbidities were also considered during decision making. Branched devices for the preservation of the IIA seem to be safe and effective in midterm follow-up while thrombosis is a rare, early complication that may be confronted with endovascular means.<sup>11</sup> In this case, the high tortuosity of the iliac artery and the conformability of the endograft used were the main factors that affected pre-operative planning.

## CONCLUSION

The progression of the aneurysmal disease may be the Achilles's heel of aneurysm repair. Proximal and distal type I en-

doleaks could be managed adequately using completely endovascular means, as in this case. A failed previous standard EVAR may demand a more complicated endovascular approach.

**Acknowledgement:** None

**Conflict of interest:** None

## REFERENCES

- 1 Dias AP, Farivar BS, Steenberge SP, Brier C, Kuramochi Y, Lyden SP, et al. Management of failed endovascular aortic aneurysm repair with explantation or fenestrated-branched endovascular aortic aneurysm repair. *J Vasc Surg.* 2018;68:1676-1687.e3.
- 2 Kouvelos GN, Oikonomou K, Antoniou GA, Verhoeven EL, Katsargyris A. A Systematic Review of Proximal Neck Dilatation After Endovascular Repair for Abdominal Aortic Aneurysm. *J Endovasc Ther.* 2017;24:59-67.
- 3 Bastos Gonçalves F, Oliveira NF, Josee van Rijn M, Ultee KH, Hoeks SE, Ten Raa S, et al. Iliac Seal Zone Dynamics and Clinical Consequences After Endovascular Aneurysm Repair. *Eur J Vasc Endovasc Surg.* 2017;53:185-92.
- 4 Kret MR, Tran K, Lee JT. Change in Aortic Neck Diameter after Endovascular Aortic Aneurysm Repair. *Ann Vasc Surg.* 2017;43:115-20.
- 5 Kapetanios D, Banafsche R, Jerkku T, Spanos K, Hoffmann U, Fiorucci B, et al. Current evidence on aortic remodeling after endovascular repair. *J Cardiovasc Surg (Torino).* 2019;60:186-90.
- 6 Wanhainen A, Verzini F, Van Herzele I, Allaire E, Bown M, Cohnert T, et al. European Society for Vascular Surgery (ESVS) 2019 Clinical Practice Guidelines on the Management of Abdominal Aorto-iliac Artery Aneurysms. *Eur J Vasc Endovasc Surg.* 2019 Dec 12. pii: S1078-5884(19)32620-6.
- 7 Chaikof EL, Dalman RL, Eskandari MK, Jackson BM, Lee WA, Mansour MA, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. 2018;67:2-77.e2
- 8 Han Y, Zhang S, Zhang J, Ji C, Eckstein HH. Outcomes of Endovascular Abdominal Aortic Aneurysm Repair in Octogenarians: Meta-analysis and Systematic Review. *Eur J Vasc Endovasc Surg.* 2017;54:454-63
- 9 Matsagkas M, Nana P, Kouvelos G, Spanos K, Bouzia A, Mpatzalexis K, et al. Initial Experience with Chimney EVAR for the Treatment of Para- and Juxta- Renal Aneurysms. *Hel J Vasc Endovasc Surg.* 2019;2:66-72
- 10 Katsargyris A, Oikonomou K, Nagel S, Giannakopoulos T, Lg Verhoeven E. Endostaples: are they the solution to graft migration and Type I endoleaks? *J Cardiovasc Surg (Torino).* 2015;56:363-8.
- 11 Schneider DB, Milner R, Heyligers JMM, Chakfé N, Matsu-mura J. Outcomes of the GORE Iliac Branch Endoprosthesis in clinical trial and real-world registry settings. *J Vasc Surg.* 2019;69:367-77.e1.